



Synthesis of mixed-linked xylans for enzyme characterization

Boos, Irene; Clausen, Mads Hartvig

Publication date:
2017

Document Version
Publisher's PDF, also known as Version of record

[Link back to DTU Orbit](#)

Citation (APA):
Boos, I., & Clausen, M. H. (2017). *Synthesis of mixed-linked xylans for enzyme characterization*. Abstract from 19th European Carbohydrate Symposium, Barcelona, Spain.

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.



Synthesis of mixed-linked xylans for enzyme characterization

Irene Boos, Mads H. Clausen

Technical University of Denmark, DTU Chemistry, Kgs. Lyngby, Denmark.

e-mail: irboo@kemi.dtu.dk

The study of plant cell wall polysaccharides and their corresponding interactions with proteins is vital to get new insights into plant development [1]. Moreover, these polysaccharides are of interest in biotechnological research, due to their use in numerous industrial applications such as food, health care and sustainable biofuel production. A major class of hemicellulose is arabinoxylan that is an important polysaccharide component of lignocellulosic biomass [2]. To underpin the full commercial exploitation of these glycan polymers, it is necessary to learn more about the enzymatic hydrolysis of arabinoxylans. This can be achieved by chemical synthesis of well-defined oligosaccharides as models for the more complex macromolecules. Moreover, the utilization of enzyme resistant substrates can support the mapping of the active site of glycosyl-hydrolases.

The talk will highlight the synthesis of mixed *O*- and *S*-linked tetraxylans (figure 1) as possible interesting candidates for the investigation and characterization of arabinoxylan degrading enzymes [3].




Figure 1. Target structures

- [1] a) R. A. Burton, M. J. Gidley, G. B. Fincher, Heterogeneity in the chemistry, structure and function of plant cell walls, *Nat. Chem. Biol.* **2010**, 6, 724.
b) K. H. Caffall, D. Mohnen, The structure, function, and biosynthesis of plant cell wall pectic polysaccharides, *Carbohydr. Res.* **2009**, 344, 1879.
- [2] B. C. Saha, Hemicellulose bioconversion, *J. Ind. Microbiol. Biotechnol.* **2003**, 30, 279.
- [3] B. Bonora, I. Boos, M. H. Clausen, Convergent strategy for the synthesis of S-linked oligoxylans, submitted to *Carbohydr. Res.*